

# **Chapter 15**



## **Multiple Integrals**

15.2 Double Integrals over genaral regions

## Definition.




Given: A function  $f$  defined on  $D$

Extend  $f$  to a rectangle containing  $D$




$$\iint_D f(x, y) \, dA = \iint_R F(x, y) \, dA$$

Region of type I.





$$D = \{(x, y) \mid a \leq x \leq b, g_1(x) \leq y \leq g_2(x)\}$$



**EXAMPLE 1** Evaluate  $\iint_D (x + 2y) dA$ , where  $D$  is the region bounded by the parabolas  $y = 2x^2$  and  $y = 1 + x^2$ .

Region of Type II.

$$D = \{(x, y) \mid c \leq y \leq d, h_1(y) \leq x \leq h_2(y)\}$$



$$\iint_D f(x, y) dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy$$

**EXAMPLE.** Evaluate  $\iint_D e^{-y^2} dA$ , where  $D$  is the region bounded by the lines  $x = 0$ ,  $y = 3$  and  $x = y$ .

**EXAMPLE.** Find the volume of the tetrahedron bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $y = 0$ , and  $z = 0$ .

**EXAMPLE 5** Evaluate the iterated integral  $\int_0^1 \int_x^1 \sin(y^2) dy dx$ .

## Properties of Double Integrals.


$$[6] \quad \iint_D (f(x, y) + g(x, y)) dA = \iint_D f(x, y) dA + \iint_D g(x, y) dA$$

$$[7] \quad \iint_D c f(x, y) dA = c \iint_D f(x, y) dA$$

$$[8] \quad \text{If } f(x, y) \geq g(x, y) \text{ on } D, \text{ then } \iint_D f(x, y) dA \geq \iint_D g(x, y) dA$$

$$[9] \quad \text{If } D = D_1 \cup D_2, \text{ with } D_1 \cap D_2 = \emptyset, \text{ then}$$


$$\iint_D f(x, y) dA = \iint_{D_1} f(x, y) dA + \iint_{D_2} f(x, y) dA$$



$$[10] \quad \text{Area}(D) = \iint_D 1 dA$$

$$[11] \quad \text{If } m \leq f(x, y) \leq M, \text{ then } m \cdot \text{Area}(D) \leq \iint_D f(x, y) dA \leq M \cdot \text{Area}(D)$$

**Example.** Find the area of the trapezoid below:



**Challenge.** Find the area of the hexagon below using properties 9 and 10:

