

## Introduction

We saw in class that the area of a surface  $S$  given by the parametric equation  $\vec{r}(u, v)$ , where  $(u, v) \in D$ , is given by the following formula

$$\text{Area}(S) = \iint_S dS = \iint_D |\vec{r}_u \times \vec{r}_v| dA,$$

where  $dA$  is the area differential adapted to  $D$  and

$$\vec{r}_u \times \vec{r}_v = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ x_u & y_u & z_u \\ x_v & y_v & z_v \end{vmatrix}.$$

In this worksheet, we will derive formulas for the area of specific types of surfaces.

## Area of Graphs

The first type of surface we will work with is the graph of a function in two variables. This is usually given by an expression of the following form:

$$z = f(x, y)$$

with  $(x, y)$  restricted to some region  $D$  in the  $xy$ -plane. A parametrization of a surface  $S$  given by the equation  $z = f(x, y)$  is

$$\vec{r}(x, y) = \langle x, y, f(x, y) \rangle,$$

where  $(x, y) \in D$ . Using this parametrization, we find that

$$\vec{r}_x \times \vec{r}_y = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 0 & f_x \\ 0 & 1 & f_y \end{vmatrix} = \langle -f_x, -f_y, 1 \rangle.$$

Hence, the surface area differential takes the following form:

$$dS = |\vec{r}_x \times \vec{r}_y| dA = \sqrt{1 + f_x^2 + f_y^2} dA.$$

The formula for area of the surface  $S$  given by the equation  $z = f(x, y)$ , where  $f$  has continuous partial derivatives, is then

$$\text{Area}(S) = \iint_D \sqrt{1 + f_x^2 + f_y^2} dA. \tag{1}$$

**PROBLEM 1.** Use Formula (1) to compute the area of the following surfaces.

a)  $z = x^2 + y^2$ , with  $x^2 + y^2 \leq 4$ .

b)  $z = \cos(x) \cos(y)$ , with  $0 \leq x \leq y \leq \pi$  (need to estimate the value of the integral).

## Area of Surfaces of Revolution

In Calculus I, you probably encountered surfaces of revolution about the  $x$ -axis, about the  $y$ -axis or maybe about any horizontal/vertical lines. The second type of surfaces we will study are the surfaces of revolution. Our focus here will be on surfaces of revolution obtained by rotating the graph  $y = f(x)$  about the  $x$ -axis.

**PROBLEM 2.** Suppose a graph of a function is given by the equation  $y = x$ , where  $0 \leq x \leq 1$ . Write or copy/paste the following link <https://www.desmos.com/3d/689fbd2154> in your favorite web browser. Then answer the following questions in order.

- Move the cursor controlling the parameter  $a$ . What is it doing?
- With the parameter  $a$  set to 1, move the cursor controlling the parameter  $b$ . What is the orange circle doing?
- The radius of the orange circle is  $f(x)$ . Use that to deduce the coordinates  $y$  and  $z$  of the points  $(x, y, z)$  on the surface you see in Desmos.

In general, a parametrization of the surface of revolution generated by the graph of  $y = f(x)$  is

$$\vec{r}(x, \theta) = \langle x, f(x) \cos(\theta), f(x) \sin(\theta) \rangle,$$

where  $a \leq x \leq b$  and  $0 \leq \theta \leq 2\pi$ . From this parametrization, we find that

$$\vec{r}_x \times \vec{r}_\theta = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & f'(x) \cos(\theta) & f'(x) \sin(\theta) \\ 0 & -f(x) \sin(\theta) & f(x) \cos(\theta) \end{vmatrix} = \langle f(x)f'(x), -f(x) \cos(\theta), -f(x) \sin(\theta) \rangle.$$

Hence,

$$dS = |\vec{r}_x \times \vec{r}_\theta| dA = |f(x)| \sqrt{1 + (f'(x))^2}$$

and so

$$\text{Area}(S) = \iint_S dS = \iint_D |f(x)| \sqrt{1 + (f'(x))^2} dA \quad (2)$$

**PROBLEM 3.** Setup the integral for the area of the following surfaces of revolution.

- The surface obtained by rotating the graph of  $y = x$ ,  $0 \leq x \leq 1$ .
- The surface obtained by rotating the graph of  $y = \sin(x)$ ,  $0 \leq x \leq \pi$ .
- The surface obtained by rotating the graph of  $y = 1 + \cos(x)$ ,  $0 \leq x \leq \pi$ .

Here is a little challenge for you. Assume that a 2D curve  $C$  lies entirely above the  $x$ -axis and does not intersect itself. Assume the curve has a parametrization  $\vec{r}(t) = \langle x(t), y(t) \rangle$ , for  $a \leq t \leq b$ .

**PROBLEM 4.** Answer the following questions.

- Find a parametrization of the surface of revolution obtained by rotating the curve  $C$  about the  $x$ -axis.
- Modify the code from Desmos to draw the surface obtained by rotating the circle with equation  $(0.5 \cos(2\pi t), 1 + 0.5 \sin(2\pi t))$ ,  $0 \leq t \leq 1$ . What is the surface?
- Find an expression of the area of the surface generated.